

***Gonepteryx farinosa* (Zeller, 1847) and *Pieris krueperi* Staudinger, 1860 confirmed for Albania, with a list of species observed in June 1995 (Lepidoptera: Papilionoidea)**

Stanislav Abadjiev & Stoyan Beshkov

Samenvatting. *Gonepteryx farinosa* (Zeller, 1847) en *Pieris krueperi* Staudinger, 1860 met zekerheid vastgesteld in Albanië, met een lijst van de waargenomen soorten in juni 1995 (Lepidoptera: Papilionoidea) Na een afwezigheidsperiode van meer dan 60 jaar kon de aanwezigheid van *Gonepteryx farinosa* (Zeller, 1847) en *Pieris krueperi* Staudinger, 1860 in Albanië bevestigd worden. Het artikel bevat een lijst van de 26 soorten dagvlinders die tussen 23 en 27 juni 1995 in Zuid-Albanië werden waargenomen.

Résumé. *Gonepteryx farinosa* (Zeller, 1847) et *Pieris krueperi* Staudinger, 1860 confirmés pour la faune d'Albanie, avec une liste des espèces observées en juin 1995 (Lepidoptera: Papilionoidea) Après une période d'absence de plus de 60 ans la présence en Albanie de *Gonepteryx farinosa* (Zeller, 1847) et *Pieris krueperi* Staudinger, 1860 pouvait être confirmée. Une liste des espèces observées entre le 23 et le 27 juin 1995 dans le sud d'Albanie est donnée.

Key words: Albania - faunistics.

Abadjiev, S.: Institute of Zoology, Bulgarian Academy of Sciences, Tsar Osvoboditel Blvd 1, BG-1000 Sofia, Bulgaria.

Beshkov, S.: 13 March Str. - 17, BG-1142 Sofia, Bulgaria.

Continuing our butterfly studies in Albania from 23rd to 27th June 1995 we visited the southern provinces of the country. On 24th June we collected in the area above Jorgucati (Province of Gjirokastra); on 25th June near Butrinti (Province of Saranda) and on 26th in the surroundings of Dhërmi (Province of Vlora). At several other sites occasional observations were done. *Gonepteryx farinosa* (Zeller, 1847) and *Pieris krueperi* Staudinger, 1860 are confirmed for Albania after a period of more than 60 years. A list of all the 26 species observed is given here.

Papilionidae

Iphiclides podalirius (Linnaeus, 1758). Observed in Butrinti and Saranda.

Papilio machaon Linnaeus, 1758. Single specimens were seen in Butrinti and Saranda.

Pieridae

Leptidea sinapis (Linnaeus, 1758). Abundant in the surroundings of Dhërmi.

Colias crocea (Fourcroy, 1785). Often seen near Jorgucati.

Gonepteryx farinosa (Zeller, 1847). A single male was taken on 24th June near Jorgucati in the same habitat as *Pieris krueperi*. Subsequently this specimen was presented to the Natural History Museum in Tirana. First reported for Albania by Rebel & Zerny (1931: 67) from Shkodra and Bojana (Province of Shkodra).

Gonepteryx cleopatra (Linnaeus, 1767). Numerous in Saranda (nectaring on *Nerium oleander*), Butrinti and Dhërmi. In the last two localities observed flying in olive plantations.

Aporia crataegi (Linnaeus, 1758). Very common on 23rd June in orchards and along roads near Pogradeci, Çerrava (Province of Pogradeci) and Mollasi (Province of Erseka).

Pieris krueperi Staudinger, 1860. Uncommon and fresh in the area above Jorgucati (Province of Gjirokastra) - 5 males, 1 female. The presence of this species on the territory of Albania is hereby confirmed. The habitat consists of an arid limestone formation, where *krueperi* flies along cliffs, commonly visiting flowers for nectaring.

Recorded by Rebel & Zerny (1931: 65) only from the Province of Librazhdi: Shkumbini Valley: Dhura, 700m, 27.VII.1917, 1 male (Bartha leg.).
Pieris mannii (Mayer, 1851). Single specimens observed in Jorgucati, Dhërmi.
Pieris rapae (Linnaeus, 1758). Common in places with secondary vegetation near Dhërmi.
Pieris napi (Linnaeus, 1758). Specimens found in Mollasi.
Pontia edusa (Fabricius, 1777). Quite common in Mollasi, Butrinti and Dhërmi.

Lycaenidae

Lampides boeticus (Linnaeus, 1767). In the coastal cliffs near Dhërmi, 1 male.
Tarucus balkanicus (Freyer, [1844]). Found along bushes of *Paliurus* near Butrinti, 2 males (in coll. Beshkov).
Pseudophilotes vicrama (Moore, 1865). Found in Dhërmi, 1 male.
Polyommatus icarus (Rottemburg, 1775). Common in Dhërmi.

Nymphalidae

Kirinia roxelana (Cramer, [1777]). Rarely found in Dhërmi, where it was flying along olive trees, sometimes resting on the trunks.
Lasiommata megera (Linnaeus, 1767). Common in rocky terrains near Jorgucati and Dhërmi.
Pyronia cecilia (Vallantin, 1894). Common and fresh in bushy places near Jorgucati, 8 males; Nivicë-Bubar (Province of Saranda), 1 male.
Hipparchia syriaca (Staudinger, 1871). Seen in rocky terrains near Jorgucati, 1 male, and along olive trees in the surroundings of Dhërmi, 2 females (in coll. Beshkov).
Hipparchia senthes (Fruhstorfer, 1908). Found near Vora (Province of Tirana), 1 female.
Melanargia larissa (Geyer, [1828]). Abundant in areas near Jorgucati, Saranda, Butrinti, Dhërmi.
Vanessa cardui (Linnaeus, 1758). Common in rocky terrains near Dhërmi.
Polygonia c-album (Linnaeus, 1758). Single specimen observed in Mollasi.
Polygonia egea (Cramer, [1775]). Only one seen near Nivicë-Bubar.
Argynnis pandora ([Denis & Schiffermüller], 1775). Single specimens seen in Dhërmi.

Reference

Rebel, H. & Zerny, H., 1931. Die Lepidopterenfauna Albaniens (Mit Berücksichtigung der Nachbargebiete). - *Denkschriften der mathem.-naturw. Klasse, Wien* 103: 37-161.